



Experiment title: Heavy atom derivative for a plant small heat shock protein	Experiment number: LS1672	
Beamline: ID14-3	Date of experiment: from: 08/06/2000 to: 09/06/2000	Date of report: 22/08/2000 <i>Received at ESRF:</i> 28 AOUT 2000
Shifts: 1	Local contact(s): Stephanie Monaco	

Names and affiliations of applicants (* indicates experimentalists):

Dr. Christine Slingsby
Dr. Rob van Montfort *

Department of Crystallography
Birkbeck College
University of London
Malet Street
London WC1E 7HX

Report:

Small heat shock proteins (sHSP) form a diverse family of proteins, which protect cells from stress by binding unfolded proteins. SHSP's have a size of 12-27 kDa, but generally form large oligomers of different sizes. The structure of the sHSP from archaean *M. Janashii* (Kim et al.) gave a first insight in such a complex, but only revealed the structure of the C-terminal domain of the two-domain sHSP monomer. To shed more light on the various functions of this protein family, we have grown crystals of a small shock protein from plant. Unfortunately, molecular replacement trials using the *M. Janashii* structure failed, due to the limited sequence homology between the *M. Janashii* structure and the plant heat shock protein. Therefore we carried out an extensive search for heavy atom derivatives and substituted the protein with selenomethionine to be able to do MAD-experiments (see also experimental reports on data collected at BM14 under proposal LS1527 and LS1672).

At the ID14-3 we have collected a data set of a crystal soaked in K₂PtCl₄. The data set was of good quality, however both difference Patterson maps and difference Fouriers using previously obtained MAD-phases showed that the heavy atom compound had not bound to the protein. Recently we have been able to solve its structure using MAD data collected at BM14 (see also experimental reports LS1527 and 1672). The structure is currently being refined.