



	Experiment title: Cellobiose dehydrogenase. BAG: Uppsala (II)	Experiment number: LS-1804
Beamline: ID14-EH4	Date of experiment: from: 4 Nov 2000 to: 6 Nov 2000	Date of report: 1 Sep 2002
Shifts: 0.2	Local contact(s): Sigrid Stuhrmann	<i>Received at ESRF:</i>
Names and affiliations of applicants (* indicates experimentalists): Christina Divne, Uppsala University, Sweden, cici@xray.bmc.uu.se * Martin Hällberg, Uppsala University, Sweden, martin-h@xray.bmc.uu.se		

Report:

The fungal flavocytochrome cellobiose dehydrogenase (CDH) participates in lignocellulose degradation with a proposed role in the early events of wood degradation. The 755-residue protein consists of a *b*-type cytochrome linked to a large FAD-binding domain which can be separated proteolytically to yield the individual proteins. CDH oxidizes cellobiose to cellobiono-1,5-lactone with the concomitant reduction of FAD. Reducing equivalents are then transferred to the cytochrome domain, and subsequently to an external electron acceptor. CDHs belong to either of two distinct groups, type-1 or type-2, that differ substantially in terms of association mode of the two domains, and in the mode of binding to the insoluble substrate (lignocellulose).

We have reported the structures of the two domains of the type-1 CDH from *Phanerochaete chrysosporium*. The structure of the CDH cytochrome revealed a for cytochromes unusual fold: a fibronectin-like β -sandwich with the protoheme bound on the surface at one face of the β -core. Moreover, a unique Met-His ligation of the *b*-type heme, *i.e.*, a near-perpendicular arrangement of the Fe ligands, was found. To investigate structural differences between the two CDH groups, the CDH cytochrome of type-2 from *Humicola insolens* will be determined. In the present experiment, a native data set was collected on $P2_12_12_1$ crystals of the *H. insolens* CDH cytochrome to 2.0 Å (R_{sym} 8.4%).