ESRF	Experiment title: High-resolution powder diffraction of a homologous series of mono-acid odd-numbered triacylglycerols.	Experiment number: CH-617
Beamline: BM16	Date of experiment: from: 11 April 1999 7:00 to: 13 April 1999 7:00	Date of report: 18 August 1999
Shifts:	Local contact(s): Andrew Fitch	Received at ESRF:

Names and affiliations of applicants (* indicates experimentalists):

Arjen van Langevelde, Wim Molleman, Eva Dova, Kees van Malssen, René Peschar, and Henk Schenk

Laboratory for Crystallography, Institute for Molecular Chemistry, Universiteit van Amsterdam, Nieuwe Achtergracht 166, 1018 WV Amsterdam, The Netherlands.

Report:

In this session a series of β - C_nC_n C_n (n = odd) triacylglycerols was measured for determination of cell parameters only (see proposal). Again, the expected cell parameters of these long-chain compounds were based upon crystallographic data of other triacylglycerols. A combination of one long (ranging from ? to ? Å) and one short cell axis (~5.5 Å) was expected and huge cell volumes. The powder diffraction patterns of $C_{13}C_{13}C_{13}$, $C_{15}C_{15}$, $C_{17}C_{17}C_{17}$ and $C_{19}C_{19}C_{19}$ were measured. From these patterns the cell parameters could be determined. From $C_{13}C_{13}C_{13}$ it was possible to solve the structure, but refinement is still not completed. With this data obtained at ESRF it must be possible to make a predictive structural model for all series members.

Since the sample of β -LOL (melting point ~15° C) was molten in the last session we decided to try it again in this session. Now, we succeeded in measuring the β polymorph of LOL and determined the cell parameters.

For structure determination three undoped poly(phenylene vinylene) compounds where measured. The patterns of these long-chain organic molecules where indexed, but no structures are solved yet.