



Experiment Report Form

The double page inside this form is to be filled in by all users or groups of users who have had access to beam time for measurements at the ESRF.

Once completed, the report should be submitted electronically to the User Office via the User Portal:
<https://www.esrf.fr/misapps/SMISWebClient/protected/welcome.do>

Deadlines for submission of Experimental Reports

Experimental reports must be submitted within the period of 3 months after the end of the experiment.

Experiment Report supporting a new proposal (“relevant report”)

If you are submitting a proposal for a new project, or to continue a project for which you have previously been allocated beam time, you must submit a report on each of your previous measurement(s):

- even on those carried out close to the proposal submission deadline (it can be a “*preliminary report*”),
- even for experiments whose scientific area is different from the scientific area of the new proposal,
- carried out on CRG beamlines.

You must then register the report(s) as “relevant report(s)” in the new application form for beam time.

Deadlines for submitting a report supporting a new proposal

- 1st March Proposal Round - **5th March**
- 10th September Proposal Round - **13th September**

The Review Committees reserve the right to reject new proposals from groups who have not reported on the use of beam time allocated previously.

Reports on experiments relating to long term projects

Proposers awarded beam time for a long term project are required to submit an interim report at the end of each year, irrespective of the number of shifts of beam time they have used.

Published papers

All users must give proper credit to ESRF staff members and proper mention to ESRF facilities which were essential for the results described in any ensuing publication. Further, they are obliged to send to the Joint ESRF/ ILL library the complete reference and the abstract of all papers appearing in print, and resulting from the use of the ESRF.

Should you wish to make more general comments on the experiment, please note them on the User Evaluation Form, and send both the Report and the Evaluation Form to the User Office.

Instructions for preparing your Report

- fill in a separate form for each project or series of measurements.
- type your report in English.
- include the experiment number to which the report refers.
- make sure that the text, tables and figures fit into the space available.
- if your work is published or is in press, you may prefer to paste in the abstract, and add full reference details. If the abstract is in a language other than English, please include an English translation.



	Experiment title: Understanding Structure and Valence in the (U, Ln⁴⁺, Ln³⁺)O₂ system (Ln³⁺ = Nd/Tm, Ln⁴⁺ = Ce)	Experiment number:
Beamline:	Date of experiment: from: 31/10/2021 to: 02/11/2022	Date of report:
Shifts:	Local contact(s): Christoph Hennig (email: hennig@esrf.fr) Volodymyr Svitlyk (email: svitlyk@esrf.fr)	<i>Received at ESRF:</i>
Names and affiliations of applicants (* indicates experimentalists): Dr Gabriel Murphy <i>Forschungszentrum Jülich GmbH, 52428, Jülich, Germany</i> Dr. Nina Huittinen Helmholtz-Zentrum Dresden-Rossendorf, 01328, Dresden, Germany		

Report:

Experiment outline:

UO₂ spent nuclear fuel possess a number of fission products such as lanthanides, Ln, which can lead to lattice distortion and loss of integrity of the fuel. The described experiment sort to obtain high-resolution synchrotron X-ray diffraction and spectroscopy (XANES) measurements under ambient conditions for the (U, Ln⁴⁺, Ln³⁺)O₂ system where Ln³⁺ = Nd/Tm and Ln⁴⁺ = Ce. The investigation was intended to provide insight into the chemical-structural effects of minor actinide occurrence in mixed oxide nuclear fuel (MOX, Pu doped) where Ln³⁺ and Ln⁴⁺ function as surrogates for minor actinides and Pu⁴⁺ respectively and consequently are pertinent in the context of spent nuclear fuel science.

Experimental outcome

High resolution structural models were developed from collected synchrotron X-ray diffraction patterns from BM20, ROBL at the ESRF which allowed the structural effects of Ln³⁺ inclusion on the known phase separation in (U_{1-x}Ce_x)O₂ between $x = 0.15$ and 0.5 to be understood. With respect to literature, this study appears to be the first high resolution investigation of the (U, Ln⁴⁺, Ln³⁺)O₂ system performed to date. The results of the investigation are being prepared for publication at time of report submission.

.

